Table 4A Cancer Incidence in Weymouth, MA CT 4223 1982-1994

| 1982-1994 | | | Total | | | | Male | | Female | | | | |
|-----------|-----|------|-------|---------|-----|------|------|---------|--------|------|-----|---------|--|
| | Obs | Exp | SIR | 95 % CI | Obs | Exp | SIR | 95 % CI | Obs | Exp | SIR | 95 % CI | |
| Bladder | 18 | 18.8 | 96 | 57 152 | 15 | 13.8 | 108 | 61 179 | 3 | 4.9 | NC | NC | |
| Brain | 3 | 7.4 | NC | NC | 1 | 4.0 | NC | NC | 2 | 3.3 | NC | NC | |
| Kidney | 9 | 11.0 | 82 | 37 155 | 5 | 6.8 | 73.8 | 24 172 | 4 | 4.2 | NC | NC | |
| Leukemia | 8 | 9.0 | 89 | 38 175 | 5 | 5.2 | 97 | 31 226 | 3 | 3.8 | NC | NC | |
| Liver | 4 | 2.3 | NC | NC | 2 | 1.7 | NC | NC | 2 | 0.6 | NC | NC | |
| Lung | 73 | 72.0 | 101 | 79 127 | 49 | 44.2 | 111 | 82 146 | 24 | 27.8 | 86 | 55 129 | |
| NHL | 17 | 16.6 | 102 | 60 164 | 9 | 8.9 | 101 | 46 192 | 8 | 7.7 | 103 | 44 203 | |
| Pancreas | 7 | 9.8 | 71 | 29 147 | 5 | 4.9 | 102 | 33 239 | 2 | 4.9 | NC | NC | |

Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

Table 4B Cancer Incidence in Weymouth, MA CT 4223

1982-1986 and 1987-1994

| 1982-1986 | | | Total | | | | | Male | | Female | | | | |
|-----------|-----|------|-------|----|------|-----|------|-------|------|--------|-----|-----|-----|---------|
| | Obs | Exp | SIR | 95 | % CI | Obs | Exp | SIR | 95 % | 6 CI | Obs | Exp | SIR | 95 % CI |
| Bladder | 6 | 7.4 | 81 | 30 | 176 | 6 | 5.5 | 109 | 39.8 | - 237 | 0 | 1.9 | NC | NC |
| Brain | 1 | 2.5 | NC | | NC | 0 | 1.4 | NC | N | С | 1 | 1.1 | NC | NC |
| Kidney | 5 | 3.3 | 151 | 49 | 351 | 3 | 2.0 | NC | No | C | 2 | 1.3 | NC | NC |
| Leukemia | 3 | 3.4 | NC | | NC | 2 | 1.9 | NC | N | С | 1 | 1.5 | NC | NC |
| Liver | 1 | 0.7 | NC | | NC | 1 | 0.5 | NC | N | С | 0 | 0.2 | NC | NC |
| Lung | 22 | 25.9 | 85 | 53 | 128 | 18 | 16.9 | 106.4 | 63 | 168 | 4 | 9.0 | NC | NC |
| NHL | 5 | 5.3 | 94 | 30 | 220 | 2 | 2.8 | NC | N | C | 3 | 2.5 | NC | NC |
| Pancreas | 2 | 3.8 | NC | | NC | 1 | 1.9 | NC | No | С | 1 | 1.9 | NC | NC |

| 1987-1994 | | | Total | | | | | Male | | Female | | | | |
|-----------|-----|------|-------|----|------|-----|------|------|----------|--------|------|-----|---------|--|
| | Obs | Exp | SIR | 95 | % CI | Obs | Exp | SIR | 95 % CI | Obs | Exp | SIR | 95 % CI | |
| Bladder | 12 | 11.3 | 106 | 55 | 185 | 9 | 8.3 | 108 | 49 205 | 3 | 3.0 | NC | NC | |
| Brain | 2 | 4.8 | NC | | NC | 1 | 2.6 | NC | NC | 1 | 2.2 | NC | NC | |
| Kidney | 4 | 7.7 | NC | | NC | 2 | 4.7 | NC | NC | 2 | 2.9 | NC | NC | |
| Leukemia | 5 | 5.6 | 89 | 29 | 209 | 3 | 3.2 | NC | NC | 2 | 2.4 | NC | NC | |
| Liver | 3 | 1.6 | NC | | NC | 1 | 1.2 | NC | NC | 2 | 0.4 | NC | NC | |
| Lung | 51 | 46.2 | 110 | 82 | 145 | 31 | 27.4 | 113 | 76.9 161 | 20 | 18.8 | 107 | 65 165 | |
| NHL | 12 | 11.3 | 106 | 55 | 185 | 7 | 6.1 | 115 | 46 237 | 5 | 5.2 | 95 | 31 223 | |
| Pancreas | 5 | 6.0 | 83 | 27 | 193 | 4 | 3.0 | NC | NC | 1 | 3.0 | NC | NC | |

Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health